

**WEIGHTED REVERSE INEQUALITIES OF JENSEN TYPE FOR
 FUNCTIONS OF SELFADJOINT OPERATORS**

S.S. DRAGOMIR

ABSTRACT. On making use of the representation in terms of the Riemann-Stieltjes integral of spectral families for selfadjoint operators in Hilbert spaces, we establish here some weighted reverse inequalities of Jensen's type for convex functions of operators. Some applications for simple functions of operators are also provided.

1. INTRODUCTION

Let $(\Omega, \mathcal{A}, \mu)$ be a measurable space consisting of a set Ω , a σ -algebra \mathcal{A} of parts of Ω and a countably additive and positive measure μ on \mathcal{A} with values in $\mathbb{R} \cup \{\infty\}$. For a μ -measurable function $w : \Omega \rightarrow \mathbb{R}$, with $w(x) \geq 0$ for μ -a.e. (almost every) $x \in \Omega$, consider the Lebesgue space

$$L_w(\Omega, \mu) := \left\{ f : \Omega \rightarrow \mathbb{R}, f \text{ is } \mu\text{-measurable and } \int_{\Omega} w(x) |f(x)| d\mu(x) < \infty \right\}.$$

For simplicity of notation we write everywhere in the sequel $\int_{\Omega} w d\mu$ instead of $\int_{\Omega} w(x) d\mu(x)$.

In order to provide a reverse of the celebrated Jensen's integral inequality for convex functions, the author obtained in 2002 [6] the following result:

Theorem 1. *Let $\Phi : [m, M] \subset \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable convex function on (m, M) and $f : \Omega \rightarrow [m, M]$ so that $\Phi \circ f, f, \Phi' \circ f, (\Phi' \circ f) f \in L_w(\Omega, \mu)$, where $w \geq 0$ μ -a.e. (almost everywhere) on Ω with $\int_{\Omega} w d\mu = 1$. Then we have the inequality:*

$$\begin{aligned} 0 &\leq \int_{\Omega} (\Phi \circ f) w d\mu - \Phi \left(\int_{\Omega} f w d\mu \right) \\ &\leq \int_{\Omega} (\Phi' \circ f) f w d\mu - \int_{\Omega} (\Phi' \circ f) w d\mu \int_{\Omega} f w d\mu \\ &\leq \frac{1}{2} [\Phi'(M) - \Phi'(m)] \int_{\Omega} w \left| f - \int_{\Omega} f w d\mu \right| d\mu. \end{aligned} \tag{1}$$

If $\mu(\Omega) < \infty$ and $\Phi \circ f, f, \Phi' \circ f, (\Phi' \circ f) f \in L(\Omega, \mu)$, then we have the inequality:

$$\begin{aligned} 0 &\leq \frac{1}{\mu(\Omega)} \int_{\Omega} (\Phi \circ f) d\mu - \Phi \left(\frac{1}{\mu(\Omega)} \int_{\Omega} f d\mu \right) \\ &\leq \frac{1}{\mu(\Omega)} \int_{\Omega} (\Phi' \circ f) f d\mu - \frac{1}{\mu(\Omega)} \int_{\Omega} (\Phi' \circ f) d\mu \frac{1}{\mu(\Omega)} \int_{\Omega} f d\mu \\ &\leq \frac{1}{2} [\Phi'(M) - \Phi'(m)] \frac{1}{\mu(\Omega)} \int_{\Omega} \left| f - \frac{1}{\mu(\Omega)} \int_{\Omega} f d\mu \right| d\mu. \end{aligned} \tag{2}$$

The following discrete inequality is of interest as well.

1991 *Mathematics Subject Classification.* Primary 26D15, 26D20; Secondary 47A63.
Key words and phrases. Jensen's inequality, Measurable functions, Lebesgue integral, Selfadjoint operators, Spectral family.

Corollary 1. Let $\Phi : [m, M] \rightarrow \mathbb{R}$ be a differentiable convex function on (m, M) . If $x_i \in [m, M]$ and $w_i \geq 0$ ($i = 1, \dots, n$) with $W_n := \sum_{i=1}^n w_i = 1$, then one has the counterpart of Jensen's weighted discrete inequality:

$$\begin{aligned} 0 &\leq \sum_{i=1}^n w_i \Phi(x_i) - \Phi\left(\sum_{i=1}^n w_i x_i\right) \\ &\leq \sum_{i=1}^n w_i \Phi'(x_i) x_i - \sum_{i=1}^n w_i \Phi'(x_i) \sum_{i=1}^n w_i x_i \\ &\leq \frac{1}{2} [\Phi'(M) - \Phi'(m)] \sum_{i=1}^n w_i \left| x_i - \sum_{j=1}^n w_j x_j \right|. \end{aligned} \quad (3)$$

Remark 1. We notice that the inequality between the first and the second term in (3) was proved in 1994 by Dragomir & Ionescu, see [24].

If $f, g : \Omega \rightarrow \mathbb{R}$ are μ -measurable functions and $f, g, fg \in L_w(\Omega, \mu)$, then we may consider the Čebyšev functional

$$T_w(f, g) := \int_{\Omega} wfgd\mu - \int_{\Omega} wfd\mu \int_{\Omega} wgd\mu. \quad (4)$$

The following result is known in the literature as the *Grüss inequality*

$$|T_w(f, g)| \leq \frac{1}{4} (\Gamma - \gamma) (\Delta - \delta), \quad (5)$$

provided

$$-\infty < \gamma \leq f(x) \leq \Gamma < \infty, \quad -\infty < \delta \leq g(x) \leq \Delta < \infty \quad (6)$$

for μ -a.e. (almost every) $x \in \Omega$.

The constant $\frac{1}{4}$ is sharp in the sense that it cannot be replaced by a smaller constant.

If we assume that $-\infty < \gamma \leq f(x) \leq \Gamma < \infty$ for μ -a.e. $x \in \Omega$, then by the Grüss inequality for $g = f$ and by the Schwarz's integral inequality, we have

$$\int_{\Omega} w \left| f - \int_{\Omega} wfd\mu \right| d\mu \leq \left[\int_{\Omega} wf^2d\mu - \left(\int_{\Omega} wfd\mu \right)^2 \right]^{\frac{1}{2}} \leq \frac{1}{2} (\Gamma - \gamma). \quad (7)$$

In what follows, we assume that $w : \Omega \rightarrow \mathbb{R}$, with $w(x) \geq 0$ for μ -a.e. $x \in \Omega$, is a μ -measurable function with $\int_{\Omega} wd\mu = 1$.

On making use of the results (1) and (7), we can state the following string of reverse inequalities:

Lemma 1. Let $\Phi : I \rightarrow \mathbb{R}$ be a differentiable convex function on the interval of real numbers I and $m, M \in \mathbb{R}$, $m < M$ with $[m, M] \subset \overset{\circ}{I}$, $\overset{\circ}{I}$ is the interior of I . If $f : \Omega \rightarrow \mathbb{R}$ is μ -measurable, satisfies the bounds

$$-\infty < m \leq f(x) \leq M < \infty \quad \text{for } \mu\text{-a.e. } x \in \Omega \quad (8)$$

and $\Phi \circ f, f, \Phi' \circ f, (\Phi' \circ f) f \in L_w(\Omega, \mu)$, then

$$\begin{aligned} 0 &\leq \int_{\Omega} (\Phi \circ f) wd\mu - \Phi\left(\int_{\Omega} f wd\mu\right) \\ &\leq \int_{\Omega} (\Phi' \circ f) f wd\mu - \int_{\Omega} (\Phi' \circ f) wd\mu \int_{\Omega} f wd\mu \\ &\leq \frac{1}{2} [\Phi'(M) - \Phi'(m)] \int_{\Omega} \left| f - \int_{\Omega} wfd\mu \right| wd\mu \end{aligned} \quad (9)$$

$$\begin{aligned} &\leq \frac{1}{2} [\Phi'(M) - \Phi'(m)] \left[\int_{\Omega} f^2 w d\mu - \left(\int_{\Omega} f w d\mu \right)^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{4} [\Phi'(M) - \Phi'(m)] (M - m). \end{aligned}$$

Remark 2. We notice that the inequality between the first, second and last term from (9) was proved in the general case of positive linear functionals in 2001 by the author in [5].

If the differentiability condition is removed, then Φ' can be replaced in the inequality (9) above with a section φ of the subdifferential $\partial\Phi$. We omit the details.

The following reverse of the Jensen's inequality holds [20], [21]:

Lemma 2. Let $\Phi : I \rightarrow \mathbb{R}$ be a convex function on the interval of real numbers I and $m, M \in \mathbb{R}$, $m < M$ with $[m, M] \subset I$. If $f : \Omega \rightarrow \mathbb{R}$ is μ -measurable, satisfies the bounds (8) and $f, \Phi \circ f \in L_w(\Omega, \mu)$, then

$$\begin{aligned} 0 &\leq \int_{\Omega} (\Phi \circ f) w d\mu - \Phi \left(\int_{\Omega} f w d\mu \right) \\ &\leq \frac{(M - \int_{\Omega} f w d\mu) (\int_{\Omega} f w d\mu - m)}{M - m} \sup_{t \in (m, M)} \Psi_{\Phi}(t; m, M) \\ &\leq \left(M - \int_{\Omega} f w d\mu \right) \left(\int_{\Omega} f w d\mu - m \right) \frac{\Phi'_-(M) - \Phi'_+(m)}{M - m} \\ &\leq \frac{1}{4} (M - m) [\Phi'_-(M) - \Phi'_+(m)], \end{aligned} \quad (10)$$

where $\Psi_{\Phi}(\cdot; m, M) : (m, M) \rightarrow \mathbb{R}$ is defined by

$$\Psi_{\Phi}(t; m, M) = \frac{\Phi(M) - \Phi(t)}{M - t} - \frac{\Phi(t) - \Phi(m)}{t - m}.$$

We also have the inequality

$$\begin{aligned} 0 &\leq \int_{\Omega} (\Phi \circ f) w d\mu - \Phi \left(\int_{\Omega} f w d\mu \right) \leq \frac{1}{4} (M - m) \Psi_{\Phi} \left(\int_{\Omega} f w d\mu; m, M \right) \\ &\leq \frac{1}{4} (M - m) [\Phi'_-(M) - \Phi'_+(m)], \end{aligned} \quad (11)$$

provided that $\int_{\Omega} f w d\mu \in (m, M)$.

We also have:

Lemma 3. With the assumptions of Lemma 2, we have the inequalities

$$\begin{aligned} 0 &\leq \int_{\Omega} w (\Phi \circ f) d\mu(x) - \Phi \left(\int_{\Omega} f w d\mu \right) \\ &\leq 2 \max \left\{ \frac{M - \int_{\Omega} f w d\mu}{M - m}, \frac{\int_{\Omega} f w d\mu - m}{M - m} \right\} \\ &\quad \times \left[\frac{\Phi(m) + \Phi(M)}{2} - \Phi \left(\frac{m + M}{2} \right) \right] \\ &\leq \frac{1}{2} \max \left\{ M - \int_{\Omega} f w d\mu, \int_{\Omega} f w d\mu - m \right\} [\Phi'_-(M) - \Phi'_+(m)]. \end{aligned} \quad (12)$$

Since

$$\frac{M - \int_{\Omega} f w d\mu}{M - m}, \frac{\int_{\Omega} f w d\mu - m}{M - m} \leq 1$$

we also have the simpler inequality

$$\begin{aligned} 0 &\leq \int_{\Omega} w (\Phi \circ f) d\mu(x) - \Phi \left(\int_{\Omega} f w d\mu \right) \\ &\leq 2 \left[\frac{\Phi(m) + \Phi(M)}{2} - \Phi \left(\frac{m + M}{2} \right) \right]. \end{aligned} \quad (13)$$

The discrete case of this inequality was obtained in [41].

For a real function $g : [m, M] \rightarrow \mathbb{R}$ and two distinct points $\alpha, \beta \in [m, M]$ we recall that the *divided difference* of g in these points is defined by

$$[\alpha, \beta; g] := \frac{g(\beta) - g(\alpha)}{\beta - \alpha}.$$

The following result holds [22]:

Lemma 4. *Let $\Phi : I \rightarrow \mathbb{R}$ be a convex function on the interval of real numbers I and $m, M \in \mathbb{R}$, $m < M$ with $[m, M] \subset I$. If $f : \Omega \rightarrow \mathbb{R}$ is μ -measurable, satisfying the bounds (8) and $f, \Phi \circ f \in L_w(\Omega, \mu)$, then by assuming that $\int_{\Omega} w f d\mu \neq m, M$, we have*

$$\begin{aligned} &\left| \int_{\Omega} \left| \Phi(f) - \Phi \left(\int_{\Omega} w f d\mu \right) \right| \operatorname{sgn} \left[f - \int_{\Omega} w f d\mu \right] w d\mu \right| \\ &\leq \int_{\Omega} (\Phi \circ f) w d\mu - \Phi \left(\int_{\Omega} w f d\mu \right) \\ &\leq \frac{1}{2} \left(\left[\int_{\Omega} w f d\mu, M; \Phi \right] - \left[m, \int_{\Omega} w f d\mu; \Phi \right] \right) D_w(f) \\ &\leq \frac{1}{2} \left(\left[\int_{\Omega} w f d\mu, M; \Phi \right] - \left[m, \int_{\Omega} w f d\mu; \Phi \right] \right) D_{w,2}(f) \\ &\leq \frac{1}{4} \left(\left[\int_{\Omega} w f d\mu, M; \Phi \right] - \left[m, \int_{\Omega} w f d\mu; \Phi \right] \right) (M - m), \end{aligned} \quad (14)$$

where

$$D_w(f) := \int_{\Omega} w \left| f - \int_{\Omega} w f d\mu \right| d\mu$$

and

$$D_{w,2}(f) := \left[\int_{\Omega} w f^2 d\mu - \left(\int_{\Omega} w f d\mu \right)^2 \right]^{\frac{1}{2}}.$$

The constant $\frac{1}{2}$ in the second inequality from (10) is best possible.

For recent results related to Jensen's inequality, see [1]-[7], [26]-[42] and the references therein.

2. INEQUALITIES FOR SELFADJOINT OPERATORS

Let A be a selfadjoint operator on the complex Hilbert space $(H, \langle \cdot, \cdot \rangle)$ with the spectrum $Sp(A)$ included in the interval $[m, M]$ for some real numbers $m < M$ and let $\{E_{\lambda}\}_{\lambda}$ be its *spectral family*. Then for any continuous function $f : [m, M] \rightarrow \mathbb{R}$, it is well known

that we have the following *spectral representation in terms of the Riemann-Stieltjes integral* (see for instance [26, p. 257]):

$$\langle f(A)x, y \rangle = \int_{m-0}^M f(\lambda) d\langle E_\lambda x, y \rangle, \quad (15)$$

and

$$\|f(A)x\|^2 = \int_{m-0}^M |f(\lambda)|^2 d\|E_\lambda x\|^2, \quad (16)$$

for any $x, y \in H$.

The function $g_{x,y}(\lambda) := \langle E_\lambda x, y \rangle$ is of *bounded variation* on the interval $[m, M]$ and $g_{x,y}(m-0) = 0$ while $g_{x,y}(M) = \langle x, y \rangle$ for any $x, y \in H$. It is also well known that $g_x(\lambda) := \langle E_\lambda x, x \rangle$ is *monotonic nondecreasing* and *right continuous* on $[m, M]$ for any $x \in H$.

The following result that provides an operator version for the Jensen inequality:

Theorem 2 (Mond-Pečarić, 1993, [35]). *Let A be a selfadjoint operator on the Hilbert space H and assume that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. If Φ is a convex function on $[m, M]$, then*

$$\Phi(\langle Ax, x \rangle) \leq \langle \Phi(A)x, x \rangle \quad (\text{MP})$$

for each $x \in H$ with $\|x\| = 1$.

As a special case of Theorem 2 we have the following Hölder-McCarthy inequality:

Theorem 3 (Hölder-McCarthy, 1967, [32]). *Let A be a selfadjoint positive operator on a Hilbert space H . Then for all $x \in H$ with $\|x\| = 1$,*

- (i) $\langle A^r x, x \rangle \geq \langle Ax, x \rangle^r$ for all $r > 1$;
- (ii) $\langle A^r x, x \rangle \leq \langle Ax, x \rangle^r$ for all $0 < r < 1$;
- (iii) If A is invertible, then $\langle A^r x, x \rangle \geq \langle Ax, x \rangle^r$ for all $r < 0$.

The following reverse for the (MP) inequality that generalizes the scalar Lah-Ribarić inequality for convex functions is well known, see for instance [25, p. 57]:

Theorem 4. *Let A be a selfadjoint operator on the Hilbert space H and assume that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. If Φ is a convex function on $[m, M]$, then*

$$\langle \Phi(A)x, x \rangle \leq \frac{M - \langle Ax, x \rangle}{M - m} \Phi(m) + \frac{\langle Ax, x \rangle - m}{M - m} \Phi(M) \quad (\text{LR})$$

for each $x \in H$ with $\|x\| = 1$.

In [23] we obtained the following weighted version of (MP) and (LR).

Theorem 5. *Let A be a selfadjoint operator on the Hilbert space H and assume that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. If $\Phi : [k, K] \subset \mathbb{R} \rightarrow \mathbb{R}$ is a continuous convex function on the interval $[k, K]$, $w : [m, M] \rightarrow [0, \infty)$ is continuous on $[m, M]$, $f : [m, M] \subset \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function on the interval $[m, M]$ and with the property that*

$$k \leq f(t) \leq K \text{ for any } t \in [m, M], \quad (17)$$

then

$$\Phi\left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \leq \frac{\langle w(A)(\Phi \circ f)(A)x, x \rangle}{\langle w(A)x, x \rangle}$$

$$\leq \frac{\left(K - \frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \Phi(k) + \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} - k\right) \Phi(K)}{K - k}, \quad (18)$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

For various particular instances of (18) that are of interest being related to Hölder-McCarthy's inequalities mentioned above, see [23].

For classical and recent result concerning inequalities for continuous functions of selfadjoint operators, see [35], [34], [36], [31], [25], [6], [10], [13], [18], [11], [15], [19], [17], [16], [14], [9], and [12].

3. FURTHER REVERSE INEQUALITIES

We have the following new results:

Theorem 6. *Let A be a selfadjoint operator on the Hilbert space H such that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. Assume that $\Phi : [k, K] \subset \mathbb{R} \rightarrow \mathbb{R}$ is a continuous convex function on the interval $[k, K]$, $w : [m, M] \rightarrow [0, \infty)$ is continuous on $[m, M]$, $f : [m, M] \subset \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function on the interval $[m, M]$ and satisfies the property (17)*

(i) *If Φ is continuously differentiable on (k, K) , then we have*

$$\begin{aligned} 0 &\leq \frac{\langle w(A)(\Phi \circ f)(A)x, x \rangle}{\langle w(A)x, x \rangle} - \Phi\left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \\ &\leq \frac{\langle (\Phi' \circ f)(A)f(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle} \\ &\quad - \frac{\langle (\Phi' \circ f)(A)w(A)x, x \rangle \langle f(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle \langle w(A)x, x \rangle} \\ &\leq \frac{1}{2} [\Phi'_-(K) - \Phi'_+(k)] \frac{\left\langle \left| f(A) - \frac{\langle f(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle} \mathbf{1}_H \right| x, x \right\rangle}{\langle w(A)x, x \rangle} \\ &\leq \frac{1}{2} [\Phi'_-(K) - \Phi'_+(k)] \left[\frac{\langle f^2(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle} - \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{4} [\Phi'_-(K) - \Phi'_+(k)] (K - k) \end{aligned} \quad (19)$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

(ii) *If we consider the function $\Psi_\Phi(\cdot; k, K) : (k, K) \rightarrow \mathbb{R}$ defined by*

$$\Psi_\Phi(t; k, K) = \frac{\Phi(K) - \Phi(t)}{K - t} - \frac{\Phi(t) - \Phi(k)}{t - k},$$

then

$$\begin{aligned} 0 &\leq \frac{\langle w(A)(\Phi \circ f)(A)x, x \rangle}{\langle w(A)x, x \rangle} - \Phi\left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \\ &\leq \frac{\left(K - \frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} - k\right)}{K - k} \sup_{t \in (k, K)} \Psi_\Phi(t; k, K) \\ &\leq \left(K - \frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} - k\right) \frac{\Phi'_-(K) - \Phi'_+(k)}{K - k} \\ &\leq \frac{1}{4} [\Phi'_-(K) - \Phi'_+(k)] (K - k) \end{aligned} \quad (20)$$

and

$$\begin{aligned}
0 &\leq \frac{\langle w(A)(\Phi \circ f)(A)x, x \rangle}{\langle w(A)x, x \rangle} - \Phi \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} \right) \\
&\leq \frac{1}{4}(K-k)\Psi_{\Phi} \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K \right) \\
&\leq \frac{1}{4}[\Phi'_-(K) - \Phi'_+(k)](K-k)
\end{aligned} \tag{21}$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

(iii) We have the inequalities

$$\begin{aligned}
0 &\leq \frac{\langle w(A)(\Phi \circ f)(A)x, x \rangle}{\langle w(A)x, x \rangle} - \Phi \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} \right) \\
&\leq 2 \max \left\{ \frac{K - \frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}}{K-k}, \frac{\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} - k}{K-k} \right\} \\
&\quad \times \left[\frac{\Phi(k) + \Phi(K)}{2} - \Phi \left(\frac{k+K}{2} \right) \right]
\end{aligned} \tag{22}$$

and

$$\begin{aligned}
0 &\leq \frac{\langle w(A)(\Phi \circ f)(A)x, x \rangle}{\langle w(A)x, x \rangle} - \Phi \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} \right) \\
&\leq 2 \left[\frac{\Phi(k) + \Phi(K)}{2} - \Phi \left(\frac{k+K}{2} \right) \right]
\end{aligned} \tag{23}$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

(iv) We also have the inequalities

$$\begin{aligned}
0 &\leq \frac{\langle w(A)(\Phi \circ f)(A)x, x \rangle}{\langle w(A)x, x \rangle} - \Phi \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} \right) \\
&\leq \frac{1}{2}\Psi_{\Phi} \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K \right) \frac{\left\langle \left| f(A) - \frac{\langle f(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle} 1_H \right| x, x \right\rangle}{\langle w(A)x, x \rangle} \\
&\leq \frac{1}{2}\Psi_{\Phi} \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K \right) \\
&\quad \times \left[\frac{\langle f^2(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle} - \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\
&\leq \frac{1}{4}\Psi_{\Phi} \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K \right) (K-k)
\end{aligned} \tag{24}$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

Proof. (i) Let $\{E_{\lambda}\}_{\lambda}$ be the spectral family of the operator A . Let $\varepsilon > 0$ and write the inequality (9) on the interval $[m - \varepsilon, M]$ and for the monotonic nondecreasing function $g(t) = \langle E_t x, x \rangle$, $x \in H$ with $\langle w(A)x, x \rangle \neq 0$, to get

$$\begin{aligned}
0 &\leq \frac{\int_{m-\varepsilon}^M (\Phi \circ f)(t) w(t) d \langle E_t x, x \rangle}{\int_{m-\varepsilon}^M w(t) d \langle E_t x, x \rangle} - \Phi \left(\frac{\int_{m-\varepsilon}^M f(t) w(t) d \langle E_t x, x \rangle}{\int_{m-\varepsilon}^M w(t) d \langle E_t x, x \rangle} \right) \\
&\leq \frac{\int_{m-\varepsilon}^M (\Phi' \circ f)(t) f(t) w(t) d \langle E_t x, x \rangle}{\int_{m-\varepsilon}^M w(t) d \langle E_t x, x \rangle}
\end{aligned} \tag{25}$$

$$\begin{aligned}
& - \frac{\int_{m-\varepsilon}^M (\Phi' \circ f)(t) w(t) d\langle E_t x, x \rangle}{\int_{m-\varepsilon}^M w(t) d\langle E_t x, x \rangle} \frac{\int_{m-\varepsilon}^M f(t) w(t) d\langle E_t x, x \rangle}{\int_{m-\varepsilon}^M w(t) d\langle E_t x, x \rangle} \\
& \leq \frac{1}{2} \frac{[\Phi'_-(K) - \Phi'_+(k)]}{\int_{m-\varepsilon}^M w(t) d\langle E_t x, x \rangle} \\
& \times \int_{m-\varepsilon}^M \left| f(t) - \frac{\int_{m-\varepsilon}^M f(s) w(s) d\langle E_s x, x \rangle}{\int_{m-\varepsilon}^M w(s) d\langle E_s x, x \rangle} \right| w(t) d\langle E_t x, x \rangle \\
& \leq \frac{1}{2} [\Phi'_-(K) - \Phi'_+(k)] \\
& \times \left[\frac{\int_{m-\varepsilon}^M f^2(t) w(t) d\langle E_t x, x \rangle}{\int_{m-\varepsilon}^M w(s) d\langle E_s x, x \rangle} - \left(\frac{\int_{m-\varepsilon}^M f(s) w(s) d\langle E_s x, x \rangle}{\int_{m-\varepsilon}^M w(s) d\langle E_s x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\
& \leq \frac{1}{4} [\Phi'_-(K) - \Phi'_+(k)] (K - k).
\end{aligned}$$

Letting $\varepsilon \rightarrow 0+$ and using the spectral representation theorem summarized in (15) we get the required inequality (19).

(ii) Follows by Lemma 2, (iii) follows by Lemma 3 while (iv) follows by Lemma 4. The details are omitted. \square

We have the following generalization and reverse for the Hölder-McCarthy inequality:

Corollary 2. *Let A be a selfadjoint operator on the Hilbert space H such that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. Assume that $w : [m, M] \rightarrow [0, \infty)$ is continuous on $[m, M]$, $f : [m, M] \subset \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function on the interval $[m, M]$ and satisfies the property (17) with $k > 0$. Assume also that $p \in (-\infty, 0) \cup (1, \infty)$.*

(i) *We have*

$$\begin{aligned}
0 & \leq \frac{\langle w(A) f^p(A) x, x \rangle}{\langle w(A) x, x \rangle} - \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right)^p \tag{26} \\
& \leq p \left[\frac{\langle f^p(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle} - \frac{\langle f^{p-1}(A) w(A) x, x \rangle \langle f(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle^2} \right] \\
& \leq \frac{1}{2} p (K^{p-1} - k^{p-1}) \frac{\left\langle \left| f(A) - \frac{\langle f(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle} 1_H \right| x, x \right\rangle}{\langle w(A) x, x \rangle} \\
& \leq \frac{1}{2} p (K^{p-1} - k^{p-1}) \left[\frac{\langle f^2(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle} - \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\
& \leq \frac{1}{4} p (K^{p-1} - k^{p-1}) (K - k)
\end{aligned}$$

for any $x \in H$ with $\langle w(A) x, x \rangle \neq 0$.

(ii) *If we consider the function $\Psi_p(\cdot; k, K) : (k, K) \rightarrow \mathbb{R}$ defined by*

$$\Psi_p(t; k, K) = \frac{K^p - t^p}{K - t} - \frac{t^p - k^p}{t - k},$$

then

$$0 \leq \frac{\langle w(A) f^p(A) x, x \rangle}{\langle w(A) x, x \rangle} - \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right)^p \tag{27}$$

$$\begin{aligned}
&\leq \frac{\left(K - \frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} - k\right)}{K - k} \sup_{t \in (k, K)} \Psi_p(t; k, K) \\
&\leq p \frac{K^{p-1} - k^{p-1}}{K - k} \left(K - \frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right) \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} - k\right) \\
&\leq \frac{1}{4}p (K^{p-1} - k^{p-1}) (K - k)
\end{aligned}$$

and

$$\begin{aligned}
0 &\leq \frac{\langle w(A)f^p(A)x, x \rangle}{\langle w(A)x, x \rangle} - \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right)^p \\
&\leq \frac{1}{4}(K - k) \Psi_p\left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K\right) \\
&\leq \frac{1}{4}p (K^{p-1} - k^{p-1}) (K - k)
\end{aligned} \tag{28}$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

(iii) We have the inequalities

$$\begin{aligned}
0 &\leq \frac{\langle w(A)f^p(A)x, x \rangle}{\langle w(A)x, x \rangle} - \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right)^p \\
&\leq 2 \max \left\{ \frac{K - \frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}}{K - k}, \frac{\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle} - k}{K - k} \right\} \\
&\quad \times \left[\frac{k^p + K^p}{2} - \left(\frac{k + K}{2}\right)^p \right]
\end{aligned} \tag{29}$$

and

$$\begin{aligned}
0 &\leq \frac{\langle w(A)f^p(A)x, x \rangle}{\langle w(A)x, x \rangle} - \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right)^p \\
&\leq 2 \left[\frac{k^p + K^p}{2} - \left(\frac{k + K}{2}\right)^p \right]
\end{aligned} \tag{30}$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

(iv) We also have the inequalities

$$\begin{aligned}
0 &\leq \frac{\langle w(A)f^p(A)x, x \rangle}{\langle w(A)x, x \rangle} - \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right)^p \\
&\leq \frac{1}{2} \Psi_p\left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K\right) \frac{\left\langle \left| f(A) - \frac{\langle f(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle} 1_H \right| x, x \right\rangle}{\langle w(A)x, x \rangle} \\
&\leq \frac{1}{2} \Psi_p\left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K\right) \\
&\quad \times \left[\frac{\langle f^2(A)w(A)x, x \rangle}{\langle w(A)x, x \rangle} - \left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}\right)^2 \right]^{\frac{1}{2}} \\
&\leq \frac{1}{4} \Psi_p\left(\frac{\langle w(A)f(A)x, x \rangle}{\langle w(A)x, x \rangle}; k, K\right) (K - k)
\end{aligned} \tag{31}$$

for any $x \in H$ with $\langle w(A)x, x \rangle \neq 0$.

If $p \in (0, 1)$, then by taking $\Phi(t) = -t^p$ we can get similar inequalities. However the details are omitted.

If we take $\Phi(t) = -\ln t$, $t > 0$ in Theorem 6 then we get the following logarithmic inequalities:

Corollary 3. *Let A be a selfadjoint operator on the Hilbert space H such that $Sp(A) \subseteq [m, M]$ for some scalars m, M with $m < M$. Assume that $w : [m, M] \rightarrow [0, \infty)$ is continuous on $[m, M]$, $f : [m, M] \subset \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function on the interval $[m, M]$ and satisfies the property (17) with $k > 0$.*

(i) *We have*

$$\begin{aligned}
0 &\leq \ln \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) - \frac{\langle w(A) \ln f(A) x, x \rangle}{\langle w(A) x, x \rangle} \\
&\leq \frac{\langle f^{-1}(A) w(A) x, x \rangle \langle f(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle^2} - 1 \\
&\leq \frac{1}{2} \frac{K - k}{kK} \left\langle \left| f(A) - \frac{\langle f(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle} 1_H \right| x, x \right\rangle \\
&\leq \frac{1}{2} \frac{K - k}{kK} \left[\frac{\langle f^2(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle} - \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\
&\leq \frac{1}{4} \frac{(K - k)^2}{kK}
\end{aligned} \tag{32}$$

for any $x \in H$ with $\langle w(A) x, x \rangle \neq 0$,

(ii) *If we consider the function $\Psi_{-\ln}(\cdot; k, K) : (k, K) \rightarrow \mathbb{R}$ defined by*

$$\Psi_{-\ln}(t; k, K) = \frac{\ln t - \ln k}{t - k} - \frac{\ln K - \ln t}{K - t},$$

then

$$\begin{aligned}
0 &\leq \ln \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) - \frac{\langle w(A) \ln f(A) x, x \rangle}{\langle w(A) x, x \rangle} \\
&\leq \frac{\left(K - \frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} - k \right)}{K - k} \sup_{t \in (k, K)} \Psi_{-\ln}(t; k, K) \\
&\leq \frac{1}{Kk} \left(K - \frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} - k \right) \leq \frac{1}{4} \frac{(K - k)^2}{kK}
\end{aligned} \tag{33}$$

and

$$\begin{aligned}
0 &\leq \ln \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) - \frac{\langle w(A) \ln f(A) x, x \rangle}{\langle w(A) x, x \rangle} \\
&\leq \frac{1}{4} (K - k) \Psi_{-\ln} \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle}; k, K \right) \leq \frac{1}{4} \frac{(K - k)^2}{kK}
\end{aligned} \tag{34}$$

for any $x \in H$ with $\langle w(A) x, x \rangle \neq 0$.

(iii) *We have the inequalities*

$$\begin{aligned}
0 &\leq \ln \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) - \frac{\langle w(A) \ln f(A) x, x \rangle}{\langle w(A) x, x \rangle} \\
&\leq 2 \max \left\{ \frac{K - \frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle}}{K - k}, \frac{\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} - k}{K - k} \right\} \ln \left(\frac{k + K}{2\sqrt{kK}} \right)
\end{aligned} \tag{35}$$

and

$$0 \leq \ln \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) - \frac{\langle w(A) \ln f(A) x, x \rangle}{\langle w(A) x, x \rangle} \leq \ln \left(\frac{k+K}{2\sqrt{kK}} \right)^2 \quad (36)$$

for any $x \in H$ with $\langle w(A) x, x \rangle \neq 0$.

(iv) We also have the inequalities

$$\begin{aligned} 0 &\leq \ln \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right) - \frac{\langle w(A) \ln f(A) x, x \rangle}{\langle w(A) x, x \rangle} \\ &\leq \frac{1}{2} \Psi_{-\ln} \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle}; k, K \right) \frac{\left\langle \left| f(A) - \frac{\langle f(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle} 1_H \right| x, x \right\rangle}{\langle w(A) x, x \rangle} \\ &\leq \frac{1}{2} \Psi_{-\ln} \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle}; k, K \right) \\ &\quad \times \left[\frac{\langle f^2(A) w(A) x, x \rangle}{\langle w(A) x, x \rangle} - \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{4} \Psi_{-\ln} \left(\frac{\langle w(A) f(A) x, x \rangle}{\langle w(A) x, x \rangle}; k, K \right) (K - k) \end{aligned} \quad (37)$$

for any $x \in H$ with $\langle w(A) x, x \rangle \neq 0$.

4. SOME EXAMPLES

If we choose $w(t) = 1$ and $f(t) = t$ with $t \in [m, M] \subset [0, \infty)$ then we get from Corollary 2 that

$$\begin{aligned} 0 &\leq \langle A^p x, x \rangle - \langle Ax, x \rangle^p \leq p \left[\langle A^p x, x \rangle - \langle A^{p-1} x, x \rangle \langle Ax, x \rangle \right] \\ &\leq \frac{1}{2} p (M^{p-1} - m^{p-1}) \langle |A - \langle Ax, x \rangle 1_H| x, x \rangle \\ &\leq \frac{1}{2} p (M^{p-1} - m^{p-1}) \left[\langle A^2 x, x \rangle - \langle Ax, x \rangle^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{4} p (M^{p-1} - m^{p-1}) (M - m), \end{aligned} \quad (38)$$

$$\begin{aligned} 0 &\leq \langle A^p x, x \rangle - \langle Ax, x \rangle^p \\ &\leq \frac{(M - \langle Ax, x \rangle) (\langle Ax, x \rangle - m)}{M - m} \sup_{t \in (m, M)} \Psi_p(t; m, M) \\ &\leq p \frac{M^{p-1} - m^{p-1}}{M - m} (M - \langle Ax, x \rangle) (\langle Ax, x \rangle - m) \\ &\leq \frac{1}{4} p (M^{p-1} - m^{p-1}) (M - m), \end{aligned} \quad (39)$$

$$\begin{aligned} 0 &\leq \langle A^p x, x \rangle - \langle Ax, x \rangle^p \leq \frac{1}{4} (M - m) \Psi_p(\langle Ax, x \rangle; m, M) \\ &\leq \frac{1}{4} p (M^{p-1} - m^{p-1}) (M - m), \end{aligned} \quad (40)$$

$$\begin{aligned} 0 &\leq \langle A^p x, x \rangle - \langle Ax, x \rangle^p \\ &\leq 2 \max \left\{ \frac{M - \langle Ax, x \rangle}{M - m}, \frac{\langle Ax, x \rangle - m}{M - m} \right\} \left[\frac{m^p + M^p}{2} - \left(\frac{m + M}{2} \right)^p \right], \end{aligned} \quad (41)$$

$$0 \leq \langle A^p x, x \rangle - \langle Ax, x \rangle^p \leq 2 \left[\frac{m^p + M^p}{2} - \left(\frac{m + M}{2} \right)^p \right] \quad (42)$$

and

$$\begin{aligned} 0 \leq \langle A^p x, x \rangle - \langle Ax, x \rangle^p &\leq \frac{1}{2} \Psi_p(\langle Ax, x \rangle; m, M) \langle |A - \langle Ax, x \rangle 1_H| x, x \rangle \\ &\leq \frac{1}{2} \Psi_p(\langle Ax, x \rangle; m, M) \left[\langle A^2 x, x \rangle - \langle Ax, x \rangle^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{4} \Psi_p(\langle Ax, x \rangle; m, M) (M - m) \end{aligned} \quad (43)$$

for any $x \in H$, $\|x\| = 1$.

If we choose $w(t) = t^q$, $q \neq 0$ and $f(t) = t$ with $t \in [m, M] \subset [0, \infty)$ then we get from Corollary 2 that

$$\begin{aligned} 0 &\leq \frac{\langle A^{p+q} x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right)^p \\ &\leq p \left[\frac{\langle A^{p+q} x, x \rangle}{\langle A^q x, x \rangle} - \frac{\langle A^{p+q-1} x, x \rangle}{\langle A^q x, x \rangle} \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right] \\ &\leq \frac{1}{2} p (M^{p-1} - m^{p-1}) \frac{\left\langle \left| A - \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} 1_H \right| x, x \right\rangle}{\langle A^q x, x \rangle} \\ &\leq \frac{1}{2} p (M^{p-1} - m^{p-1}) \left[\frac{\langle A^{q+2} x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{4} p (M^{p-1} - m^{p-1}) (M - m), \end{aligned} \quad (44)$$

$$\begin{aligned} 0 &\leq \frac{\langle A^{p+q} x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right)^p \\ &\leq \frac{\left(M - \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} - m \right)}{M - m} \sup_{t \in (m, M)} \Psi_p(t; m, M) \\ &\leq p \frac{M^{p-1} - m^{p-1}}{M - m} \left(M - \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} - m \right) \\ &\leq \frac{1}{4} p (M^{p-1} - m^{p-1}) (M - m), \end{aligned} \quad (45)$$

$$\begin{aligned} 0 &\leq \frac{\langle A^{p+q} x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right)^p \leq \frac{1}{4} (M - m) \Psi_p \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle}; m, M \right) \\ &\leq \frac{1}{4} p (M^{p-1} - m^{p-1}) (M - m), \end{aligned} \quad (46)$$

$$0 \leq \frac{\langle A^{p+q} x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right)^p \quad (47)$$

$$\begin{aligned} &\leq 2 \max \left\{ \frac{M - \frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle}}{M - m}, \frac{\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle} - m}{M - m} \right\} \left[\frac{m^p + M^p}{2} - \left(\frac{m + M}{2} \right)^p \right], \\ &0 \leq \frac{\langle A^{p+q}x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle} \right)^p \leq 2 \left[\frac{m^p + M^p}{2} - \left(\frac{m + M}{2} \right)^p \right] \end{aligned} \quad (48)$$

and

$$\begin{aligned} &0 \leq \frac{\langle A^{p+q}x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle} \right)^p \quad (49) \\ &\leq \frac{1}{2} \Psi_p \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle}; m, M \right) \frac{\left\langle \left| A - \frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle} 1_H \right| x, x \right\rangle}{\langle A^q x, x \rangle} \\ &\leq \frac{1}{2} \Psi_p \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle}; m, M \right) \left[\frac{\langle A^{q+2}x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{4} \Psi_p \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle}; m, M \right) (M - m) \end{aligned}$$

for any $x \in H \setminus \{0\}$.

If we choose $w(t) = 1$ and $f(t) = t$ with $t \in [m, M] \subset [0, \infty)$ then we get from Corollary 3 that

$$\begin{aligned} &0 \leq \ln \langle Ax, x \rangle - \langle \ln Ax, x \rangle \leq \langle A^{-1}x, x \rangle \langle Ax, x \rangle - 1 \quad (50) \\ &\leq \frac{1}{2} \frac{M - m}{mM} \langle |A - \langle Ax, x \rangle 1_H| x, x \rangle \\ &\leq \frac{1}{2} \frac{M - m}{mM} \left[\langle A^2x, x \rangle - \langle Ax, x \rangle^2 \right]^{\frac{1}{2}} \leq \frac{1}{4} \frac{(M - m)^2}{mM}, \end{aligned}$$

$$\begin{aligned} &0 \leq \ln \langle Ax, x \rangle - \langle \ln Ax, x \rangle \quad (51) \\ &\leq \frac{(M - \langle Ax, x \rangle) (\langle Ax, x \rangle - m)}{M - m} \sup_{t \in (m, M)} \Psi_{-\ln}(t; m, M) \\ &\leq \frac{1}{Mm} (M - \langle Ax, x \rangle) (\langle Ax, x \rangle - m) \leq \frac{1}{4} \frac{(M - m)^2}{mM}, \end{aligned}$$

$$\begin{aligned} &0 \leq \ln \langle Ax, x \rangle - \langle \ln Ax, x \rangle \leq \frac{1}{4} (M - m) \Psi_{-\ln}(\langle Ax, x \rangle; m, M) \quad (52) \\ &\leq \frac{1}{4} \frac{(M - m)^2}{mM}, \end{aligned}$$

$$\begin{aligned} &0 \leq \ln \langle Ax, x \rangle - \langle \ln Ax, x \rangle \quad (53) \\ &\leq 2 \max \left\{ \frac{M - \langle Ax, x \rangle}{M - m}, \frac{\langle Ax, x \rangle - m}{M - m} \right\} \ln \left(\frac{m + M}{2\sqrt{mM}} \right), \end{aligned}$$

$$0 \leq \ln \langle Ax, x \rangle - \langle \ln Ax, x \rangle \leq \ln \left(\frac{m + M}{2\sqrt{mM}} \right)^2 \quad (54)$$

and

$$0 \leq \ln \langle Ax, x \rangle - \langle \ln Ax, x \rangle \quad (55)$$

$$\begin{aligned}
&\leq \frac{1}{2} \Psi_{-\ln} (\langle Ax, x \rangle; m, M) \langle |f(A) - \langle Ax, x \rangle 1_H| x, x \rangle \\
&\leq \frac{1}{2} \Psi_{-\ln} (\langle Ax, x \rangle; m, M) \left[\langle A^2 x, x \rangle - \langle Ax, x \rangle^2 \right]^{\frac{1}{2}} \\
&\leq \frac{1}{4} \Psi_{-\ln} (\langle Ax, x \rangle; m, M) (M - m)
\end{aligned}$$

for any $x \in H$ with $\|x\| = 1$.

If we choose $w(t) = t^q$, $q \neq 0$ and $f(t) = t$ with $t \in [m, M] \subset [0, \infty)$ then we get from Corollary 3 that

$$0 \leq \ln \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) - \frac{\langle A^q \ln Ax, x \rangle}{\langle A^q x, x \rangle} \quad (56)$$

$$\begin{aligned}
&\leq \frac{\langle A^{q-1} x, x \rangle \langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle \langle A^q x, x \rangle} - 1 \leq \frac{1}{2} \frac{K - k}{kK} \left\langle \left| A - \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} 1_H \right| x, x \right\rangle \\
&\leq \frac{1}{2} \frac{K - k}{kK} \left[\frac{\langle A^{q+2} x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \leq \frac{1}{4} \frac{(K - k)^2}{kK},
\end{aligned}$$

$$0 \leq \ln \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) - \frac{\langle A^q \ln Ax, x \rangle}{\langle A^q x, x \rangle} \quad (57)$$

$$\begin{aligned}
&\leq \frac{\left(K - \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} - k \right)}{K - k} \sup_{t \in (k, K)} \Psi_{-\ln}(t; k, K) \\
&\leq \frac{1}{Kk} \left(K - \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} - k \right) \leq \frac{1}{4} \frac{(K - k)^2}{kK},
\end{aligned}$$

$$0 \leq \ln \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) - \frac{\langle A^q \ln Ax, x \rangle}{\langle A^q x, x \rangle} \quad (58)$$

$$\leq \frac{1}{4} (K - k) \Psi_{-\ln} \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle}; k, K \right) \leq \frac{1}{4} \frac{(K - k)^2}{kK},$$

$$0 \leq \ln \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) - \frac{\langle A^q \ln Ax, x \rangle}{\langle A^q x, x \rangle} \quad (59)$$

$$\leq 2 \max \left\{ \frac{K - \frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle}}{K - k}, \frac{\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} - k}{K - k} \right\} \ln \left(\frac{k + K}{2\sqrt{kK}} \right),$$

$$0 \leq \ln \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) - \frac{\langle A^q \ln Ax, x \rangle}{\langle A^q x, x \rangle} \leq \ln \left(\frac{k + K}{2\sqrt{kK}} \right)^2, \quad (60)$$

and

$$0 \leq \ln \left(\frac{\langle A^{q+1} x, x \rangle}{\langle A^q x, x \rangle} \right) - \frac{\langle A^q \ln Ax, x \rangle}{\langle A^q x, x \rangle} \quad (61)$$

$$\begin{aligned}
&\leq \frac{1}{2} \Psi_{-\ln} \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle}; k, K \right) \frac{\left\langle \left| A - \frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle} 1_H \right| x, x \right\rangle}{\langle A^q x, x \rangle} \\
&\leq \frac{1}{2} \Psi_{-\ln} \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle}; k, K \right) \left[\frac{\langle A^{q+2}x, x \rangle}{\langle A^q x, x \rangle} - \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle} \right)^2 \right]^{\frac{1}{2}} \\
&\leq \frac{1}{4} \Psi_{-\ln} \left(\frac{\langle A^{q+1}x, x \rangle}{\langle A^q x, x \rangle}; k, K \right) (K - k)
\end{aligned}$$

for any $x \in H \setminus \{0\}$.

REFERENCES

- [1] Abramovich, S., *Convexity, subadditivity and generalized Jensen's inequality*, Ann. Funct. Anal. **4** (2013), no. 2, 183–194.
- [2] Aldaz, J.M., *A measure-theoretic version of the Dragomir-Jensen inequality*, Proc. Amer. Math. Soc. **140** (2012), no. 7, 2391–2399.
- [3] Cirtoaje, V., *The best lower bound for Jensen's inequality with three fixed ordered variables*, Banach J. Math. Anal. **7** (2013), no. 1, 116–131.
- [4] Dragomir, S.S., *A converse result for Jensen's discrete inequality via Grüss' inequality and applications in information theory*, An. Univ. Oradea Fasc. Mat. **7** (1999/2000), 178–189.
- [5] Dragomir, S.S., *On a reverse of Jensen's inequality for isotonic linear functionals*, J. Ineq. Pure & Appl. Math. **2**(2001), No. 3, Article 36.
- [6] Dragomir, S.S., *A Grüss type inequality for isotonic linear functionals and applications*, Demonstratio Math. **36** (2003), no. 3, 551–562. Preprint RGMIA Res. Rep. Coll. **5**(2002), Supplement, Art. 12. [Online [http://rgmia.org/v5\(E\).php](http://rgmia.org/v5(E).php)].
- [7] Dragomir, S.S., *Bounds for the normalized Jensen functional*, Bull. Austral. Math. Soc. **74** (3) (2006), 471–476.
- [8] Dragomir, S.S., *Bounds for the deviation of a function from the chord generated by its extremities*, Bull. Aust. Math. Soc. **78** (2008), no. 2, 225–248.
- [9] Dragomir, S.S., *Some inequalities for convex functions of selfadjoint operators in Hilbert spaces*, Filomat **23** (2009), No. 3, 81–92. Preprint RGMIA Res. Rep. Coll., **11**(e) (2008), Art. 10.
- [10] Dragomir, S.S., *Grüss' type inequalities for functions of selfadjoint operators in Hilbert spaces*, Preprint, RGMIA Res. Rep. Coll., **11**(e) (2008), Art. 11. [ONLINE: [http://rgmia.org/v11\(E\).php](http://rgmia.org/v11(E).php)].
- [11] Dragomir, S.S., *Some reverses of the Jensen inequality for functions of selfadjoint operators in Hilbert spaces*, J. Ineq. & Appl. Vol. **2010**, Article ID 496821. Preprint RGMIA Res. Rep. Coll. **11**(e) (2008), Art. 15. [ONLINE: [http://rgmia.org/v11\(E\).php](http://rgmia.org/v11(E).php)].
- [12] Dragomir, S.S., *Some Jensen's type inequalities for twice differentiable functions of selfadjoint operators in Hilbert spaces*, Filomat **23** (2009), No. 3, 211–222. Preprint RGMIA Res. Rep. Coll. **11**(e) (2008), Art. 13.
- [13] Dragomir, S.S., *Some new Grüss' type inequalities for functions of selfadjoint operators in Hilbert spaces*, Sarajevo J. Math. **6** (18), (2010), No. 1, 89–107. Preprint RGMIA Res. Rep. Coll. **11**(e) (2008), Art. 12. [ONLINE: [http://rgmia.org/v11\(E\).php](http://rgmia.org/v11(E).php)].
- [14] Dragomir, S.S., *Some Slater's type inequalities for convex functions of selfadjoint operators in Hilbert spaces*, Rev. Un. Mat. Argentina, **52** (2011), No.1, 109–120. Preprint RGMIA Res. Rep. Coll. **11**(e) (2008), Art. 7.
- [15] Dragomir, S.S., *Some Jensen's type inequalities for log-convex functions of selfadjoint operators in Hilbert spaces*, Bull. Malays. Math. Sci. Soc. **34**(2011), No. 3. Preprint RGMIA Res. Rep. Coll. **13**(2010), Sup. Art. 2.
- [16] Dragomir, S.S., *Hermite-Hadamard's type inequalities for operator convex functions*, Appl. Math. Comp. **218** (2011), 766–772. Preprint RGMIA Res. Rep. Coll. **13** (2010), No. 1, Art. 7.
- [17] Dragomir, S.S., *Hermite-Hadamard's type inequalities for convex functions of selfadjoint operators in Hilbert spaces*, Preprint RGMIA Res. Rep. Coll. **13** (2010), No. 2, Art. 1.
- [18] Dragomir, S.S., *New bounds for the Čebyšev functional of two functions of selfadjoint operators in Hilbert spaces*, Filomat **24** (2010), No. 2, 27–39.

- [19] Dragomir, S.S., *New Jensen's type inequalities for differentiable log-convex functions of selfadjoint operators in Hilbert spaces*, Sarajevo J. Math. **19** (2011), No. 1, 67-80. Preprint RGMIA Res. Rep. Coll. **13** (2010), Sup. Art. 2.
- [20] Dragomir, S.S., *Reverses of the Jensen inequality in terms of first derivative and applications*, Acta Math. Vietnam. **38** (2013), no. 3, 429–446
- [21] Dragomir, S.S., *Some reverses of the Jensen inequality with applications*, Bull. Aust. Math. Soc. **87** (2013), no. 2, 177–194.
- [22] Dragomir, S.S., *A Refinement and a divided difference reverse of Jensen's inequality with applications*, accepted Rev. Colomb. Mate. Preprint RGMIA Res. Rep Coll. **14** (2011), Art 74. [Online <http://rgmia.org/papers/v14/v14a74.pdf>].
- [23] Dragomir, S.S., *Jensen type weighted inequalities for functions of selfadjoint and unitary operators*, Ital. J. Pure Appl. Math. No. **32** (2014), 247–264.
- [24] Dragomir, S.S. and Ionescu, N.M., *Some converse of Jensen's inequality and applications*, Rev. Anal. Numér. Théor. Approx. **23** (1994), no. 1, 71–78.
- [25] Furuta, T., Mičić Hot, J., Pečarić, J. and Seo, Y., *Mond-Pečarić Method in Operator Inequalities. Inequalities for Bounded Selfadjoint Operators on a Hilbert Space*, Element, Zagreb, 2005.
- [26] Helmsberg, G., *Introduction to Spectral Theory in Hilbert Space*, John Wiley, New York, 1969.
- [27] Horváth, L., *A new refinement of discrete Jensen's inequality depending on parameters*, J. Inequal. Appl. **2013**, 2013:551, 16 pp.
- [28] Adil, K.M., Khalid, S. and Pečarić, J., *Improvement of Jensen's inequality in terms of Gâteaux derivatives for convex functions in linear spaces with applications*, Kyungpook Math. J. **52** (2012), no. 4, 495–511.
- [29] Mihai, M.V., *Jensen's inequality for mixed convex functions of two real variables*, Acta Univ. Apulensis Math. Inform. No. **34** (2013), 179–183.
- [30] Kian, M., *Operator Jensen inequality for superquadratic functions*, Linear Algebra Appl. **456** (2014), 82–87.
- [31] Matković, A., Pečarić, J. and Perić, I., *A variant of Jensen's inequality of Mercer's type for operators with applications*, Linear Algebra Appl. **418** (2006), no. 2-3, 551–564.
- [32] McCarthy, C.A., c_p , Israel J. Math. **5**(1967), 249–271.
- [33] Mičić, J., Seo, Y., Takahasi, S.-E. and Tominaga, M., *Inequalities of Furuta and Mond-Pečarić*, Math. Ineq. Appl. **2** (1999), 83–111.
- [34] Mond, B. and Pečarić, J., *On some operator inequalities*, Indian J. Math. **35** (1993), 221–232.
- [35] Mond, B. and Pečarić, J., *Convex inequalities in Hilbert space*, Houston J. Math. **19** (1993), 405–420.
- [36] Mond, B. and Pečarić, J., *Classical inequalities for matrix functions*, Utilitas Math. **46** (1994), 155–166.
- [37] Niculescu, C.P., *An extension of Chebyshev's inequality and its connection with Jensen's inequality*, J. Inequal. Appl. **6** (2001), no. 4, 451–462.
- [38] Pavić, Z., *The applications of functional variants of Jensen's inequality*, J. Funct. Spaces Appl. **2013**, Art. ID 194830, 5 pp.
- [39] Popovici, F. and Spiridon, C.-I., *The Jensen inequality for (M, N) -convex functions*, An. Univ. Craiova Ser. Mat. Inform. **38** (2011), no. 4, 63–66.
- [40] Sharma, R., *On Jensen's inequality for positive linear functionals*, Int. J. Math. Sci. Eng. Appl. **5** (2011), no. 5, 263–271.
- [41] Simić, S., *On a global upper bound for Jensen's inequality*, J. Math. Anal. Appl. **343** (2008), 414–419.
- [42] Zabandan, G. and Kiliçman, A., *A new version of Jensen's inequality and related results*, J. Inequal. Appl. **2012**, 2012:238, 7 pp.

MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE
 VICTORIA UNIVERSITY, PO BOX 14428
 MELBOURNE CITY, MC 8001, AUSTRALIA.

SCHOOL OF COMPUTATIONAL & APPLIED MATHEMATICS
 UNIVERSITY OF THE WITWATERSRAND
 PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA
E-mail address: sever.dragomir@vu.edu.au
URL: <http://rgmia.org/dragomir>